

# Arctic warming: nonlinear impacts of sea-ice and glacier melt on seabird foraging

DAVID GRÉMILLET<sup>1,2</sup>, JÉRÔME FORT<sup>3</sup>, FRANÇOISE AMÉLINEAU<sup>1</sup>, ELENA ZAKHAROVA<sup>4,5</sup>, TANGILE BOT<sup>1</sup>, ENRIC SALA<sup>6,7</sup> and MARIA GAVRILO<sup>8</sup>

<sup>1</sup>CEFE UMR 5175, CNRS – Université de Montpellier – Université Paul-Valéry Montpellier – EPHE, 1919 route de Mende, 34293 Montpellier, France, <sup>2</sup>FitzPatrick Institute, DST/NRF Excellence Centre at the University of Cape Town, Rondebosch 7701, South Africa, <sup>3</sup>LIENSs UMR 7266, CNRS – Université La Rochelle, 2 rue Olympe de Gouges, 17000 La Rochelle, France, <sup>4</sup>LEGOS UMR5566, CNRS – CNES – IRD – Université Toulouse III) 14 Avenue Edouard Belin, 31400 Toulouse, France, <sup>5</sup>State Oceanography Institute, ul. Beringa 38, 119397 St. Petersburg, Russia, <sup>6</sup>National Geographic Society, Washington, DC, USA, <sup>7</sup>Centre d'Estudis Avançats de Blanes (CEAB-CSIC), Girona, Spain, <sup>8</sup>National Park Russian Arctic, Sovetskikh Kosmonavtov Avenue, 57, 163000 Archangelsk, Russia

## Abstract

Arctic climate change has profound impacts on the cryosphere, notably via shrinking sea-ice cover and retreating glaciers, and it is essential to evaluate and forecast the ecological consequences of such changes. We studied zooplankton-feeding little auks (*Alle alle*), a key sentinel species of the Arctic, at their northernmost breeding site in Franz-Josef Land (80°N), Russian Arctic. We tested the hypothesis that little auks still benefit from pristine arctic environmental conditions in this remote area. To this end, we analysed remote sensing data on sea-ice and coastal glacier dynamics collected in our study area across 1979–2013. Further, we recorded little auk foraging behaviour using miniature electronic tags attached to the birds in the summer of 2013, and compared it with similar data collected at three localities across the Atlantic Arctic. We also compared current and historical data on Franz-Josef Land little auk diet, morphometrics and chick growth curves. Our analyses reveal that summer sea-ice retreated markedly during the last decade, leaving the Franz-Josef Land archipelago virtually sea-ice free each summer since 2005. This had a profound impact on little auk foraging, which lost their sea-ice-associated prey. Concomitantly, large coastal glaciers retreated rapidly, releasing large volumes of melt water. Zooplankton is stunned by cold and osmotic shock at the boundary between glacier melt and coastal waters, creating new foraging hotspots for little auks. Birds therefore switched from foraging at distant ice-edge localities, to highly profitable feeding at glacier melt-water fronts within <5 km of their breeding site. Through this behavioural plasticity, little auks maintained their chick growth rates, but showed a 4% decrease in adult body mass. Our study demonstrates that arctic cryosphere changes may have antagonistic ecological consequences on coastal trophic flow. Such nonlinear responses complicate modelling exercises of current and future polar ecosystem dynamics.

**Keywords:** arctic climate change, arctic marine fauna, behavioural plasticity, cryosphere, historical data, little auk

Received 7 August 2014 and accepted 14 October 2014

## Introduction

Global warming is twice as marked in the Arctic than at lower latitudes, and this has a rapid and profound impact on the cryosphere. For instance, the Intergovernmental Panel on Climate Change Fifth Assessment Report (IPCC, 2013) predicts a sea-ice-free Arctic in summer from the 2030s, or towards the end of 21st century, depending on scenarios. The Arctic being 70% marine, a reduction in sea-ice extent and thickness radically modifies its seascape, with major impacts on atmospheric and ocean circulation, biogeochemical cycles and ecosystem functioning (AMAP, 2012). Endemic arctic species, such as the emblematic

polar bear (*Ursus maritimus*), are predicted to go extinct, and to be replaced by species from lower latitudes (review in Gilg *et al.*, 2012). Yet arctic warming also leads to pronounced glacier and permafrost melt (IPCC, 2013), which strongly modify terrestrial and coastal landscapes, and associated ecological dynamics. Determining and predicting the overall impacts of cryosphere changes is therefore a key research objective (AMAP, 2012), particularly at the interface between terrestrial and marine ecosystems (Post *et al.*, 2013). Beyond polar bears, a limited number of sea-ice-associated arctic species function as sentinel species, whose ecological responses to the consequences of climate change deserve special attention. This is the case of the little auk (*Alle alle*), for a number of reasons: it is the most abundant seabird in the Atlantic

Correspondence: David Grémillet, tel. + 33 467 613 210, e-mail: david.gremillet@cefe.cnrs.fr

Arctic, with over 40 million individuals, and therefore plays a major ecological role within marine ecosystems (Karnovsky & Hunt, 2002; Karnovsky *et al.*, 2003). It is a specialized zooplankton consumer and participates in a short food chain with strong responses to sea-ice presence/absence and to sea-surface temperature (Grémillet *et al.*, 2012; Jakubas *et al.*, 2012). It is also the smallest high arctic, diving seabird, and operates at the upper limit of its energetic capabilities, something that strongly enhances its sensitivity to environmental change (Harding *et al.*, 2009a). Current knowledge shows that ocean warming and a vanishing cryosphere have a negative impact on little auks, because they preferentially feed on large, lipid-rich zooplankton (copepods) associated with cold water in the marginal ice zone (Karnovsky *et al.*, 2010). It is therefore predicted that little auk colonies situated in warmer areas will be more impacted than those from areas that are less exposed to global warming. Yet ocean warming progresses rapidly in the North Atlantic, and it seems appropriate to test the impact of cryosphere changes on little auks at their northernmost breeding site, which is situated in Franz-Josef Land (Russian State Federal Refuge), at 80°N. This extremely remote area is often regarded as an 'arctic Eden', still shielded from the consequences of global warming, with 85% of its land mass covered by glaciers and extensive, year-round, sea-ice cover surrounding islands of the archipelago. It is home for over a million seabirds, including an estimated 500 000 little auks (Anker-Nilssen, 2000), as well as large proportions of the world populations of further endemic arctic species (e.g. polar bear, walrus *Odobenus rosmarus rosmarus*, ivory gull *Pagophila eburnea* and bowhead whale *Balaena mysticetus*). Previous investigations showed that little auks mainly respond to marine environmental change through marked plasticity in their foraging behaviour and diet (Grémillet *et al.*, 2012). We therefore investigated the foraging ecology of little auks from Franz-Josef Land using miniaturized temperature–depth recorders (TDR) and dietary analyses. Adult body measurements and chick growth rates were also recorded. These fitness indicators were compared with data collected at the same study site 20 years before, and with little auk foraging behaviour from three other breeding sites across the Atlantic Arctic. Seabird data were interpreted in the context of Franz-Josef Land sea-ice and glacier dynamics, which we analysed over the past 30 years using satellite remote sensing. Specifically, we tested the hypothesis that little auks from Franz-Josef Land benefit from conditions of the pre-industrial age, with a foraging effort predicted to be lower than at other, more southern breeding sites in the Arctic.

## Materials and methods

The study site was situated in Tikhaya Bay, Hooker Island, Franz-Josef Land (80°20.293'N; 52°49.062'E), at a little auk colony facing coastal waters and in the vicinity of the 7.2 km<sup>2</sup> Sedov glacier (Fig. 1), during National Geographic's Pristine Seas 2013 Franz-Josef Land expedition. Little auks were studied in July–August 2013 during the chick-rearing phase, within the framework of a scientific programme approved by the board of the National Park Russian Arctic and the Ministry of Education and Science of the Russian Federation (Permit N°71 issued on 03/06/2013). This phase was chosen because birds are accessible, have their highest foraging effort for self-maintenance and chick-feeding, and are therefore assumed to be most susceptible to environmental change. Four different sets of birds/nests were studied synoptically within the same colony. First, 15 birds were caught in the vicinity of their nest and equipped with TDRs (G5; CEFAS Technologies, Lowestoft, UK; 31 mm length, 8 mm diameter; 2.7 g, corresponding to 1.7% of lightest handled little auk body mass, or DST micro-TD; StarOddi, Iceland; 25.4 mm length, 8.3 mm diameter; 3.3 g, corresponding to 2% of the lightest handled little auk body mass), which were attached to ventral feathers using waterproof Tesa<sup>®</sup> tape. TDRs were left on the birds for 3–6 days, upon which birds were recaptured and tags downloaded. TDRs recorded hydrostatic pressure and temperature every 4 s for the entire period (G5), or for 24 h starting on the day following deployment (DST micro-TD). Recorded temperature signals allow discrimination of periods spent at the nest, in flight and in the water and were therefore used to calculate the number of foraging trips per 24 h, foraging trip duration and flight time per trip and per 24 h (for full details on analyses see Harding *et al.*, 2009b). Pressure recordings were analysed with MULTITRACE (Jensen Software Systems, Laboe, Germany), to determine (per 24 h) the number of dives, average and maximum dive depth, average and maximum dive duration, and the per cent time spent underwater. Foraging parameters of little auks from Franz-Josef Land were compared with those of birds from East Greenland and from two locations on Svalbard, which were studied using the same methodology (data from Grémillet *et al.*, 2012). This biotelemetry study was complemented by direct observations of feeding little auks, assessed on three occasions from a boat in August 2013. During these 2-h sessions, birds were observed from <100 m on a calm sea using 12 × 42 binoculars, and positions recorded using a hand-held GPS. These at-sea observations were compared with similar surveys conducted in the summer of 2001. Second, breeding little auks carry undamaged zooplankton prey back to the nest in a sublingual pouch. Twenty birds carrying such food loads were also caught, and complete food loads stored in 70% ethanol. Prey identification was performed to the lowest possible taxonomic level under a binocular microscope. Third, we marked a set of 18 little auk nests and determined chick growth (body mass to 1 g, wing and tarsus length to 1 mm) every second day throughout the chick-rearing phase (1–26 August 2013). Finally, all birds caught for TDR deployments and diet sampling were weighed to the nearest 2 g and measured (wing and tarsus) to the nearest 1 mm. This sample was complemented with adult breeding birds caught synoptically within the same area for other purposes.

Diet, adult morphometrics and chick growth rates were compared with similar data collected by (Stempniewicz *et al.*, 1996) and (Weslawski *et al.*, 1994) within the same area in 1991–1993. We only had access to average body mass and body size measurements for 1991–1993 and therefore could not calculate/compare individual body condition indexes for the two periods.

Mean monthly sea-ice concentrations (percentage of ocean surface covered by sea-ice) were determined for July and August, across 1979–2013, within 100 km of the study colony (i.e. the maximum foraging range of breeding little auks; Anker-Nilssen, 2000; Jakubas *et al.*, 2013) using daily EUMETSAT OSI SAF ice concentration satellite products for 1979–2009. This consisted of reprocessed SMMR and SSM/I instruments data with a spatial resolution of 12.5 km (Eastwood *et al.*, 2010). AMSR-E instrument measurements with 6.25 km spatial resolution (Spreen *et al.*, 2008) were used as a complement for the 2002–2013 time period, also allowing cross-checking of the two data sets across 2002–2009.


Dynamics of the Sedov glacier were studied using high-resolution (60–15 m) Landsat images. For the 1985–2013 period, 13 images were used, and most of them acquired during July–August, the period of highest ablation, except for 1998 (April) and 2006 (September). Images were digitized, and we determined the glacier terminus position for each year. Yearly glacier progression/retreat was then calculated using 2001 as an arbitrary reference.

## Results


Summer sea-ice cover (Fig. 2a) showed strong variability during 1979–2013. Until 2003, the mean monthly sea-ice concentration within 100 km of the little auk colony varied between 40% and 70% and between 15% and 55% in July and August, respectively. Since 2003–2004, the archipelago has been almost ice free in August, because sea-ice tended to melt earlier each year. Only small ice-filled areas occurred at the north and north-east periphery of the archipelago, partly advected by winds and currents.

Coastal glaciers also retreated markedly in the study area (Fig. 2b). Since the middle of 1990s, the Sedov glacier retreated progressively, and from the year of its maximal surface area in 1994 to 2012, it has been shrinking by 0.64 km<sup>2</sup>. This corresponds to a 9% reduction of its 1994 surface (7.2 km<sup>2</sup>; 0.50% decrease per year). To verify that these observations are robust, we compared this dynamic with that of the neighbouring Yuria glacier (Fig. 1). For this glacier, the total 1985–2012 retreat of the terminus position was even higher, with an 18% decrease of its initial 12.5 km<sup>2</sup> area (0.66% per year) and a major decline between 2002 and 2012.

All 15 TDRs deployed were recovered. One data file was corrupted, and we therefore present data for 14



**Fig. 1** (a) Study area on Hooker Island, Franz-Josef Land (from a 2006 Landsat image), showing the study colony (red dot) facing Tikahaya Bay, local glaciers (white), coastal waters (blue), and approximate position of the coastal front between coastal water and glacier melt water in August 2013 (dotted line, see Data S1). (b) Situation of Franz-Josef Land (red dot) in the high Arctic.



**Fig. 2** (a) Interannual variability of August and July mean sea-ice concentration within 100 km of the little auk colony of Tikhaya Bay, from the two satellite products (SSMR + SSM/I across 1979–2009 and AMSR-E across 2002–2013). (b) Variation of the extent of the Sedov glacier ( $\text{km}^2$ ) relative to its 2001 extent.


birds (Data S1). Little auks foraged extensively, performing an average of 6 feeding trips and 361 dives per day. Dives were usually short (36 s) and shallow (6.2 m), but birds occasionally dived  $>1$  min to  $>20$  m depth. Diving effort was comparable to that of little auks diving in three other regions of the Arctic (Fig. 3a, b, see also Data S2). However, and most surprisingly, flight time per foraging trip was at least one order of magnitude lower in little auks from Franz-Josef Land than in birds from three other arctic regions (Fig. 3c, d). Such short flight times (usually  $<4$  min) between breeding colony and feeding area indicated that birds did not feed offshore, but extremely close to the colony. We confirmed this through direct at-sea observations of little auks: birds were seen between foraging dives, positioned along the front separating glacier melt from coastal waters (Fig. 1 and Data S3). Birds were only present at this front, on the sea side where water was much clearer than in glacial melt water. Conversely, at-sea surveys conducted in the summer of 2001 showed that little auks foraged in the marginal ice zone, northwest of the Franz-Josef archipelago (Data S4).

Little auk diet was largely dominated by copepods, which were present in all samples and represented 95.7% of all prey items (Table 1). These copepods were mainly *Calanus glacialis* and *C. hyperboreus*, two cold-water-associated species. This diet differed markedly

from data collected in 1991–1993 at the same study site (Weslawski *et al.*, 1994). Those showed that at that time little auks not only fed on copepods, but also to 13% on *Apherusa glacialis*, a sea-ice-associated amphipod which was present in the food of 49% of all little auks sampled in 1991–1993. This particular prey was absent from all but one of our 2013 samples (i.e. a proportion of 0.5% of all prey and 5% occurrence, see also Data S5). Growth curves of little auk chicks measured in August 1992 (Stempniewicz *et al.*, 1996) and in August 2013 (our study) were identical (Fig. 4). Adult body measurements ( $n = 57$  in 1992 and  $n = 108$  in 2013) showed that birds were 4% lighter in 2013 ( $194.1 \pm 11.9$  g) than in 1992 ( $202.3 \pm 12.5$  g;  $t = 4.07$ ,  $P < 0.001$ ), but that other measurements were not significantly different between the two periods (Wing length:  $133.3 \pm 3.7$  mm in 1992 and  $133.0 \pm 4.0$  mm in 2013. Tarsus length:  $23.1 \pm 0.8$  mm in 1992 and  $23.5 \pm 1.0$  mm in 2013).

## Discussion

Our results allow two important conclusions: (1) the consequences of global warming are strongly modifying coastal marine ecosystems even in Franz-Josef Land, an extremely remote, high-Arctic locality. (2) Disappearance of summer sea-ice from the archipelago and pronounced coastal glacier melt have antagonistic effects on little auk foraging behaviour, leading to a



**Fig. 3** Foraging characteristics (mean  $\pm$  SD) of little auks from four breeding colonies in the Atlantic Arctic, according to the latitude of the different sites: 70°N: Kap Hoegh, East Greenland (six birds); 77°N: Hornsund, Southern Spitsbergen (11 birds); 78°N: Kongsfjorden, Northern Spitsbergen (seven birds); 80°N: Hooker Island, Franz-Josef Land (14 birds). Data from Franz-Josef Land are from this study, others from Grémillet *et al.* (2012). Picture credits: Manfred Enstipp and David Grémillet.


nonlinear response to climate change in this endemic, ecologically highly important arctic species (Fig. 5). Our work is strongly multidisciplinary, and features the first comprehensive investigation of little auk foraging ecology in relation to sea-ice and coastal glacier melt.

Studying the overall impacts of cryosphere changes on arctic biodiversity has been identified as a major scientific challenge by the Arctic Council and its working groups (e.g. AMAP, 2012), as well as by international research initiatives such as the Belmont Forum. Specifically, there is currently a lack of studies that integrate terrestrial and marine processes and investigate the impacts of arctic climate change in coastal areas (Post *et al.*, 2013). Further, the IPCC (IPCC, 2013) and the Belmont Forum have stressed the importance of identifying nonlinear responses of organisms and populations to global change. Our research fits tightly to these objectives and builds upon existing, yet limited, work

on sea-ice-associated marine predators in the context of climate change (Gaston *et al.*, 2005; Eamer *et al.*, 2013). In a global perspective, little auks appear as the arctic ecological equivalent to Adélie penguins (*Pygoscelis adeliae*). For this species, extensive Antarctic research has demonstrated the importance of the marginal ice zone for efficient foraging on krill (*Euphausia* sp.) and fish, whereby birds require specific, intermediate sea-ice concentrations for an optimal provisioning efficiency (Emmerson & Southwell, 2008; Ballard *et al.*, 2012). Interestingly, recent investigations also demonstrated nonlinear responses of Adélie penguins to climate change, whereby their plastic phenotypic response to variable sea-ice concentration was perturbed by extreme climatic events (Lescroël *et al.*, 2014). Further, our results are also in line with recent studies and syntheses from other arctic regions, which stressed the importance of glacier outflow into coastal ecosystems,

**Table 1** Comparison of little auk diet on Hooker Island, Franz-Josef Land, in 2013 ( $n = 20$ , our study) and in 1991–1993 [ $n = 47$ , (Weslawski *et al.*, 1994)]. Note the importance of *Calanus* copepods (bold) and quasi disappearance of ice-associated *Apherusa glacialis* and *Gammarus wilkitzkii* (bold) in 2013

	1991–1993		2013	
	Relative abundance (%)	Occurrence frequency (%)	Relative abundance (%)	Occurrence frequency (%)
<b><i>Calanus</i> spp.</b>	<b>84.44</b>	<b>49.00</b>	<b>95.73</b>	<b>100.00</b>
<i>Paraeuchaeta</i> spp.	0.00	0.00	0.30	35.00
<i>Metridia longa</i>	0.00	0.00	0.42	20.00
<b><i>Apherusa glacialis</i></b>	<b>13.15</b>	<b>49.00</b>	<b>0.51</b>	<b>5.00</b>
<i>Onisimus</i> spp.	0.10	6.00	0.03	5.00
<i>Themisto</i> spp.	1.07	64.00	1.15	60.00
<i>Decapods larvae</i>	0.10	9.00	0.73	50.00
<b><i>Gammarus wilkitzkii</i></b>	<b>0.22</b>	<b>15.00</b>	<b>0.00</b>	<b>0.00</b>
<i>Thysanoessa</i> spp.	0.52	6.00	0.00	0.00
<i>Mysis oculata</i>	0.12	11.00	0.00	0.00
Fishes	0.12	9.00	0.36	40.00
<i>Ostracoda</i> spp.	0.12	2.00	0.00	0.00
<i>Acanthostepheia</i> spp.	0.02	2.00	0.00	0.00
<i>Chaetognaths</i>	0.00	0.00	0.01	5.00
Undetermined	0.00	0.00	0.77	40.00




**Fig. 4** Little auk chick growth curves from Franz-Josef Land measured in 1992 (black circles; Stempniewicz *et al.*, 1996) and in 2013 (open circles,  $n = 18$  chicks; this study). Error bars show standard deviations of the mean. Absence of error bars after 20 days is due to sample sizes of one for each age group. Picture credit: Ann Harding.

and the ecological consequences of a climate-induced increase of such outflow. In Alaska, for instance, researchers demonstrated that enhanced glacier discharge generates trophic cascades in arctic fjord ecosystems (Arimitsu *et al.*, 2012), whereby melt-water fronts attract pelagic fish suitable for seabird foraging (Renner *et al.*, 2012). For the Svalbard region, Lydersen *et al.* (2014) identified similar dynamics as those demonstrated by our study, with sea-ice rapidly disappearing as a resting and foraging habitat for a series of seabirds and marine mammals, and glacier melt water enhancing primary and secondary productivity in coastal ecosystems, to the benefit of some members of the marine predator guild. Their important conclusion was that this effect might only be temporary, as some of the coastal

glaciers might retreat from the coastline if arctic warming persists. The little auk population of Franz-Josef Land is therefore predicted to benefit from profitable foraging conditions close to shore as long as coastal glaciers are melting. According to glacier melt recorded over the last decades (0.5–0.6% of total surface per year, see Results), and assuming a linear process, a complete melt of the Sedov glacier would be completed in approximately 180 years. However, the glacier might have retreated sufficiently far inland to stop influencing coastal ecological dynamics decades before.

Karnovsky *et al.* (2010) predicted that 40% of all major little auk colonies would disappear from the Atlantic Arctic by the end of the 21st century. Their prediction, which was based on zooplankton thermal niche modelling under the influence of increasing sea-surface temperature, now has to be revisited in the context of arctic coastal glacier melt and little auk behavioural plasticity. Little auks were so far considered as an archetype of endemic, specialized arctic predator bound to disappear from a warming Arctic. Actually, they show surprising flexibility, allowing them to maintain fitness levels (Grémillet *et al.*, 2012; this study). Work conducted at little auk breeding sites across the Arctic clearly demonstrates the importance of cold water and associated large, lipid-rich copepods for their profitable foraging (Kwasniewski *et al.*, 2010; Karnovsky *et al.*, 2011; Frandsen *et al.*, 2014), as well as their close association with the marginal ice zone (Jakubas *et al.*, 2013). However, our study demonstrates that glacier melt water can also generate optimal foraging conditions for this species, as a refuge coastal habitat in



**Fig. 5** Hypothetical nonlinear impacts of sea-ice disappearance and coastal glacier melt onto the performance (foraging profitability and fitness) of ice-associated seabirds. Slopes are arbitrary. Note that sea-ice disappearance is initially predicted to have a negative impact on seabirds (1), until glacier discharge is sufficiently strong to create a new coastal foraging habitat (2); current situation in Franz-Josef Land). Reduction of glacier discharge in the absence of sea-ice is then predicted to further diminish seabird performance (3), but foraging plasticity might allow birds to switch to another feeding mode and stabilize/improve their performance (dotted line). Picture credits: NASA and David Grémillet.

the absence of sea-ice. Higher variability in diving effort for little auks from Franz-Josef Land compared to other localities (Fig. 3a, b) nonetheless suggests lower predictability in food availability at glacier melt-water fronts. Further, it is unclear to us why little auks make such profitable use of this new foraging habitat in Franz-Josef Land, and not at other arctic localities. This may be due to the geography of the archipelago, with a very high number of small islands covered by glaciers. Franz-Josef Land therefore has many more glacier fronts exposed to coastal waters than Svalbard, for instance. This presumption calls for further studies assessing the importance of coastal glacier frontal zones to foraging little auks at the meta-population level.

There is currently a huge demand for predicting the fate of arctic biodiversity exposed to ongoing climate change. At the species level, this is achieved by building habitat models linking distribution ranges with a suite of abiotic and biotic environmental variables (Elith & Leathwick, 2009). This environmental envelope is then extrapolated to generate predicted future distributions (e.g. Péron *et al.*, 2012; Beaugrand *et al.*, 2014). In this context, and as clearly demonstrated by our study, it is essential to identify and implement key environmental drivers. Ignoring some of these factors, and neglecting nonlinear, plastic behavioural responses of the species involved (Fig. 5) may lead to erroneous predictions (Barnagaud *et al.*, 2012; La Sorte & Jetz, 2012; Schurr *et al.*, 2012).

## Acknowledgements

This study was conducted in the specially protected area of the Franz-Josef Land State Federal Refuge, within the framework of the joint international expedition of National Park Russian Arctic and a National Geographic 'Pristine Seas Expedition FJL 2013'. The expedition was funded by National Geographic, Blancpain and Davidoff Cool Water. DG and JF are funded by CNRS and the European Union (Marie Curie CIG; Project 631203 to JF), and by the French Polar Institute (ADACLIM program No. 388). We are grateful to the Russian Geographical Society for its support and to the Russian Arctic National Park for research permits. We warmly thank the international 'Pristine Seas' team and all persons present on board *Polaris* during the expedition. We are also immensely grateful to all Russian colleagues present at Tikhaya Bay station in August 2013 for their hospitality, in particular to our bodyguard Vladimir Kretov for endless patience during fieldwork. We thank Craig R. White for advice on allometry, and Delphine Bonnet and Rumsais Blatrix for their assistance throughout the zooplankton analyses.

## References

- AMAP (2012) *Arctic Climate Issues 2011: Changes in Arctic Snow, Water, Ice and Permafrost*. SWIPA 2011 Overview Report. Arctic Monitoring and Assessment Programme (AMAP), Oslo.
- Anker-Nilssen T (2000) The status of marine birds breeding in the Barents Sea region. *Norsk Polarinstitutt Rapport* No. 113. pp. 131–136.
- Arimitsu ML, Piatt JF, Madison EN, Conaway JS, Hillgruber N (2012) Oceanographic gradients and seabird prey community dynamics in glacial fjords. *Fisheries Oceanography*, **21**, 148–169.
- Ballard G, Jongsomjit D, Veloz SD, Ainley DG (2012) Coexistence of mesopredators in an intact polar ocean ecosystem: the basis for defining a Ross Sea marine protected area. *Biological Conservation*, **156**, 72–82.
- Barnagaud J-Y, Devicor V, Jiguet F, Barbet-Massin M, Le Viol I, Archaux F (2012) Relating habitat and climatic niches in birds. *PLoS One*, **7**, e32819.
- Beaugrand G, Goberville E, Lyczak C, Kirby RR (2014) Marine biological shifts and climate. *Proceedings of the Royal Society B-Biological Sciences*, **281**, UNSP 20133350.
- Eamer J, Donaldson G, Gaston T *et al.* (2013) *Life Linked to Ice. A Guide to Sea-Ice Associated Biodiversity in this Time of Rapid Change*. Conservation of Arctic Flora and Fauna (CAFF), Akureyri, Iceland.
- Eastwood S, Larsen KR, Lavergne T, Nielsen E, Tonboe R (2010) Global Sea Ice Concentration Reprocessing – Product User Manual, *EUMETSAT OSI SAF report*. SAF/OSI/CDOP/met.no/TEC/MA/138, v1.1.
- Elith J, Leathwick JR (2009) Species distribution models: ecological explanation and prediction across space and time. *Annual Review of Ecology Evolution and Systematics*, **40**, 677–697.
- Emmerson L, Southwell C (2008) Sea ice cover and its influence on Adelie Penguin reproductive performance. *Ecology*, **89**, 2096–2102.
- Frandsen MS, Fort J, Rigét FF, Galatius A, Mosbech A (2014) Composition of chick meals from one of the main little auk (*Alle alle*) breeding colonies in Northwest Greenland. *Polar Biology*, **37**, 1055–1060.
- Gaston AJ, Gilchrist HG, Hipfner J (2005) Climate change, ice conditions and reproduction in an Arctic nesting marine bird: Brunnich's guillemot (*Uria lomvia* L.). *Journal of Animal Ecology*, **74**, 832–841.
- Gilg O, Kovacs KM, Aars J *et al.* (2012) Climate change and the ecology and evolution of Arctic vertebrates. *Annals of the New York Academy of Sciences*, **1249**, 166–190.
- Grémillet D, Welcker J, Karnovsky NJ *et al.* (2012) Little auks buffer the impact of current Arctic climate change. *Marine Ecology Progress Series*, **454**, 197–206.
- Harding AMA, Kitaysky AS, Hall ME *et al.* (2009a) Flexibility in the parental effort of an Arctic-breeding seabird. *Functional Ecology*, **23**, 348–358.
- Harding AMA, Egevang C, Walkusz W, Merkel F, Blanc S, Grémillet D (2009b) Estimating prey capture rates of a planktivorous seabird, the little auk (*Alle alle*), using diet, diving behaviour, and energy consumption. *Polar Biology*, **32**, 785–796.
- IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. (eds Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM). Cambridge University Press, Cambridge, UK and New York, NY.
- Jakubas D, Iliszko L, Wojczulanis-Jakubas K, Stempniewicz L (2012) Foraging by little auks in the distant marginal sea ice zone during the chick-rearing period. *Polar Biology*, **35**, 73–81.
- Jakubas D, Trudnowska E, Wojczulanis-Jakubas K *et al.* (2013) Foraging closer to the colony leads to faster growth in little auks. *Marine Ecology Progress Series*, **489**, 263–278.
- Karnovsky NJ, Hunt GL (2002) Estimation of carbon flux to dovekeys (*Alle alle*) in the North Water. *Deep Sea Research Part II: Topical Studies in Oceanography*, **49**, 5117–5130.
- Karnovsky NJ, Kwasiński S, Weslawski JM, Walkusz W, Beszczynska-Møller A (2003) Foraging behavior of little auks in a heterogeneous environment. *Marine Ecology Progress Series*, **253**, 289–303.
- Karnovsky N, Harding A, Walkusz W *et al.* (2010) Foraging distribution of little auks (*Alle alle*) across the Greenland Sea: implications of present and future Arctic climate change. *Marine Ecology Progress Series*, **415**, 283–293.
- Karnovsky NJ, Brown ZW, Welcker J *et al.* (2011) Inter-colony comparison of diving behavior of an Arctic top predator: implications for warming in the Greenland Sea. *Marine Ecology Progress Series*, **440**, 229–240.
- Kwasniewski S, Gluchowska M, Jakubas D *et al.* (2010) The impact of different hydrographic conditions and zooplankton communities on provisioning Little Auks along the West coast of Spitsbergen. *Progress in Oceanography*, **87**, 72–82.
- La Sorte FA, Jetz W (2012) Tracking of climatic niche boundaries under recent climate change. *Journal of Animal Ecology*, **81**, 914–925.
- Lescoërl A, Ballard G, Grémillet D, Authier M, Ainley DG (2014) Antarctic climate change: extreme events disrupt plastic phenotypic response in Adelie Penguins. *PLoS One*, **9**, e85291.
- Lydersen C, Assmy P, Falk-Petersen S *et al.* (2014) The importance of tidewater glaciers for marine mammals and seabirds in Svalbard, Norway. *Journal of Marine Systems*, **129**, 452–471.
- Péron C, Weimerskirch H, Bost C-A (2012) Projected poleward shift of king penguins' (*Aptenodytes patagonicus*) foraging range at the Crozet Islands, southern Indian Ocean. *Proceedings of the Royal Society B-Biological Sciences*, **279**, 2515–2523.
- Post E, Bhatt US, Bitz CM *et al.* (2013) Ecological consequences of sea-ice decline. *Science*, **341**, 519–524.
- Renner M, Arimitsu ML, Piatt JF (2012) Structure of marine predator and prey communities along environmental gradients in a glaciated fjord. *Canadian Journal of Fisheries and Aquatic Sciences*, **69**, 2029–2045.
- Schurr FM, Pagel J, Cabral JS *et al.* (2012) How to understand species' niches and range dynamics: a demographic research agenda for biogeography. *Journal of Biogeography*, **39**, 2146–2162.
- Spreen G, Kaleschke L, Heygster G (2008) Sea ice remote sensing using AMSR-E 89-GHz channels. *Journal of Geophysical Research: Oceans*, **113**, C02S03.
- Stempniewicz L, Skakuj M, Iliszko L (1996) The little auk *Alle alle polaris* of Franz Josef Land: a comparison with Svalbard *Alle alle* populations. *Polar Research*, **15**, 1–10.
- Weslawski JM, Stempniewicz L, Galaktionov K (1994) Summer diet of seabirds from the Frans Josef Land archipelago, Russian Arctic. *Polar Research*, **13**, 173–181.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Data S1.** Foraging characteristics of 14 chick-rearing little auks from Tikhaya Bay, Franz-Josef Land in August 2013. Recordings for 1–6 days (38 bird-days in total).

**Data S2.** Considering allometry in inter-colony comparisons of little auk foraging effort.

**Data S3.** Little auk foraging hotspot at the interface between glacier melt and coastal waters.

**Data S4.** Sea-ice concentration and little auk at-sea sightings in August 2001.

**Data S5.** Diet of 20 chick-rearing little auks from Tikhaya Bay, Franz-Josef Land in August 2013.